

# ADVICE TO HOUSEHOLDERS

YOUR PROPERTY HAS BEEN TREATED  
WITH RODENTICIDE TODAY BY:

COMPANY/AUTHORITY

Date:

Operators name:

Product:

*(Operator to insert product name)*

This section **MUST** be  
completed by the:  
Pest Control Operative  
**OR** the Relevent Authority

## CONTAINS THE FOLLOWING ACTIVE INGREDIENT:

**BRODIFACOUM**

**DIFENACOUM**

**WARFARIN**

*(Operator to tick appropriate box)*

## THIS PRODUCT CONTAINS AN ANTICOAGULANT.

You are advised that this rodenticide is approved for use in domestic premises, e.g. houses and flats, under the Control of Pesticides Regulations 1986. However, your co-operation is requested in adopting the following precautions to further minimise any risks to occupants and pets.

## PRECAUTIONS

**AVOID ALL CONTACT BY MOUTH.**

**PREVENT ACCESS TO BAIT** by children, birds and non-target animals (particularly dogs, cats, pigs and poultry).

**IN CASE OF ACCIDENT, contact a doctor or veterinarian and show this leaflet.**

## DO NOT MOVE OR TAMPER WITH BAIT.

### Advice to Physicians

If you think that bait has been eaten – The product is an indirect anticoagulant. Phytomenadione, Vitamin K1, is antidotal. Determine prothrombin times not less than eighteen hours after consumption. If elevated, administer Vitamin K1 in divided doses. Continue until prothrombin times normalise. After withdrawal of the antidote, continue determination of prothrombin time for two weeks and resume treatment if elevation occurs in that time. NB. Vitamin K3 is not effective.

For comprehensive medical advice on the treatment of poisoning, contact the nearest Poisons Information Centre or Sorex Limited.



Sorex Limited, St. Michael's Industrial Estate, Widnes, Cheshire WA8 8TJ  
Telephone: 0151 420 7151 Fax: 0151 495 1163 [www.sorex.com/pco](http://www.sorex.com/pco)

Declaration: Sorex accepts no liability if any amendments are made to this leaflet.